

CUET UG 2023 Chemistry Question Paper PDF (Jun 05, 2023) (Shift 1)

- Q1.** How much calcium will be obtained when 4 Faradays of electricity is passed through molten CaCl_2 ?
 (a) 20 g
 (b) 40 g
 (c) 80 g
 (d) 160 g
- Q2.** 0.01 M aqueous solutions of following salts were prepared. Their conductivity values were measured at 25°C . Which of the following will possess highest conductivity at 25°C ?
 (a) LiCl
 (b) RbCl
 (c) NaCl
 (d) CsCl
- Q3.** Which of the following amines can be prepared by Gabriel synthesis?
 (A) Isobutyl amine
 (B) 2-Phenyl ethylamine
 (C) N-Methyl benzyl amine
 (D) Aniline
 Choose the **correct** answer from the options given below:
 (a) (A), (B) only
 (b) (C), (D) only
 (c) (A), (C) only
 (d) (B), (D) only
- Q4.** The decomposition of H_2O_2 in alkaline medium, in presence of I^- ions occurs as follows:

$$2\text{H}_2\text{O}_2 \xrightarrow{\text{I}^-, \text{alkaline medium}} 2\text{H}_2\text{O} + \text{O}_2$$

 If the reaction occurs in the following two steps:
 Step 1: $\text{H}_2\text{O}_2 + \text{I}^- \xrightarrow{\text{Slow}} \text{H}_2\text{O} + \text{IO}^-$
 Step 2: $\text{H}_2\text{O}_2 + \text{IO}^- \xrightarrow{\text{Fast}} \text{H}_2\text{O} + \text{I}^- + \text{O}_2$
 The correct rate law expression for the reaction is:
 (a) $r = k[\text{H}_2\text{O}_2]^2 [\text{I}^-]$
 (b) $r = k[\text{H}_2\text{O}_2] [\text{I}^-]$
 (c) $r = k[\text{H}_2\text{O}_2][\text{IO}^-]$
 (d) $r = k[\text{H}_2\text{O}_2]^2 [\text{I}^-][\text{OH}^-]$
- Q5.** Identify the given name reaction:

$$\text{R}-\text{NH}_2 + \text{CHCl}_3 + 3\text{KOH} \xrightarrow{\text{Heat}} \text{R}-\text{NC} + 3\text{KCl} + 3\text{H}_2\text{O}$$

 (a) Sandmeyer's reaction
 (b) Carbylamine reaction
 (c) Hoffman's bromamide reaction
 (d) Hinsberg's reaction
- Q6.** A compound is formed by 2 elements P and Q. Atoms of element Q (as anions) makes ccp and those of P (as cations) fill all tetrahedral voids and also occupy half of the octahedral voids. What is the correct formula of this compound?
 (a) P_2Q_5
 (b) P_5Q_2
 (c) P_2Q_3
 (d) P_3Q_2
- Q7.** Arrange the following compounds in the order of their increasing boiling point:
 Pentan-1-ol, n-butane, pentanal, ethoxy ethane
 (a) n-butane < ethoxy ethane < pentanal < pentan-1-ol
 (b) n-butane < pentanal < ethoxy ethane < pentan-1-ol
 (c) n-butane < pentan-1-ol < ethoxy ethane < pentanal
 (d) pentan-1-ol < ethoxy ethane < pentanal < n-butane
- Q8.** The rate law for a reaction between substance P & Q is given by $\text{Rate} = k[\text{P}]^x[\text{Q}]^y$. If concentration of P is doubled and concentration of Q is halved, the ratio of the new rate to the earlier rate of reaction will be:
 (a) $x + y$
 (b) $x - y$
 (c) $2^x - y$
 (d) $\frac{1}{2^{x-y}}$
- Q9.** For the reactions $5\text{Br}^-(\text{aq}) + \text{BrO}_3^-(\text{aq}) + 6\text{H}^+(\text{aq}) \rightarrow 3\text{Br}_2(\text{aq}) + 3\text{H}_2\text{O}(\text{l})$. The rate of the reaction is:
 (a) one-fifth of rate of disappearance of Br^- ions
 (b) one-fifth of rate of formation of Br_2
 (c) one-third of rate of disappearance of BrO_3^- ions
 (d) one-fourth of rate of formation of H_2O
- Q10.** We have four aqueous solution labelled as:
 (A) 0.1 M NaCl
 (B) 0.01 M NaCl
 (C) 0.01 M BaCl_2
 (D) 0.01 M Sucrose
 Choose the **correct** increasing order of Van't Hoff factor from the options given below:
 (a) $i_C < i_A < i_B < i_D$
 (b) $i_D < i_C < i_B < i_A$
 (c) $i_D < i_A = i_B < i_C$
 (d) $i_C < i_A = i_B < i_D$
- Q11.** Match List - I with List - II.
- | | List - I | List - II |
|-----|------------------------------------|--------------------|
| (A) | $\text{S}_{\text{N}}1$ reaction | (I) Vic-dibromide |
| (B) | Bromination of alkenes | (II) Gem-dibromide |
| (C) | Alkylidene halides | (III) Racemisation |
| (D) | Dehydrohalogenation of haloalkanes | (IV) Saytzeff rule |
- Choose the **correct** answer from the options given below:
 (a) (A)-(IV), (B)-(II), (C)-(I), (D)-(III)
 (b) (A)-(II), (B)-(II), (C)-(I), (D)-(IV)
 (c) (A)-(IV), (B)-(I), (C)-(II), (D)-(III)
 (d) (A)-(III), (B)-(I), (C)-(II), (D)-(IV)

Q12. Which of the following statements about gold (Au) element are correct?

- (A) Gold occurs in nature in native state.
- (B) Gold dissolves in aqua-regia.
- (C) Gold is extracted by leaching the metal with dilute NaCN.

(D) Leaching is followed by replacement reaction. Choose the correct answer from the options given below:

- (a) (B), (C), (D) only
- (b) (A), (C), (D) only
- (c) (A), (B), (C) only
- (d) (A), (B), (C) and (D)

Q13. Methyl bromide reacts with silver nitrite to form:

- (a) Nitromethane
- (b) Methylnitrite
- (c) Ethanenitrite
- (d) Grignard reagent

Q14. The volume of mixing (ΔV) is not zero for which of the following solutions?

- (a) Chloroethane and Bromoethane
- (b) Benzene and Toluene
- (c) Ethanol and Acetone
- (d) n-Hexane and n-Heptane






Q15. When dilute hydrochloric acid is added to sodium chromate solution the colour changes to:

- (a) yellow
- (b) colourless
- (c) green
- (d) orange

Q16. The complex having lowest Δ_0 value among the given complexes is:

- (a) $[\text{Co}(\text{NH}_3)_6]^{3+}$
- (b) $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$
- (c) $[\text{Co}(\text{CN})_6]^{3-}$
- (d) $[\text{CrCl}_6]^{3-}$

Q17. Which one is the correct Zwitter ion of the following protein?



Q18. The actual density of a crystal of NaCl is observed to be lower than the expected value. It is caused due to the presence of:

- (a) Interstitial Defect

- (b) Frenkel Defect
- (c) Schottky Defect
- (d) Impurity Defect

Q19. The pair(s) of co-ordination complexes/ions exhibiting the same kind of isomerism is / are:

- (A) $[\text{Cr}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ and $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$
- (B) $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$ and $[\text{Pt}(\text{NH}_3)_2(\text{H}_2\text{O})\text{Cl}]^+$
- (C) $[\text{CoBr}_2\text{Cl}_2]^{-2}$ and $[\text{PtBr}_2\text{Cl}_2]^{2-}$
- (D) $[\text{Pt}(\text{NH}_3)_3(\text{NO}_3)]\text{Cl}$ and $[\text{Pt}(\text{NH}_3)_3\text{Cl}]\text{Br}$

Choose the **correct** answer from the options given below:

- (a) (A) and (C) only
- (b) (B) and (D) only
- (c) (A) and (D) only
- (d) (C) and (D) only

Q20. Match List - I with List - II.

	List - I		List - II
(A)	Polyvinyl chloride	(I)	Phenol formaldehyde
(B)	Urea-formaldehyde	(II)	Ethyleneglycol and phthalic acid
(C)	Glyptal	(III)	Vinyl chloride
(D)	Bakelite	(IV)	Urea formaldehyde

Choose the **correct** answer from the options given below:

- (a) (A)-(III), (B)-(IV), (C)-(II), (D)-(I)
- (b) (A)-(III), (B)-(IV), (C)-(I), (D)-(II)
- (c) (A)-(III), (B)-(II), (C)-(I), (D)-(IV)
- (d) (A)-(III), (B)-(II), (C)-(IV), (D)-(I)

Q21. 0.01 molal aqueous solutions of strong electrolytes NaCl, KCl, MgSO_4 and K_2SO_4 are prepared. Which of the following statements is correct regarding their osmotic pressure? (Assume complete dissociation of solute particles)

- (a) All of them exhibit equal osmotic pressure
- (b) Osmotic pressure of all of them are different
- (c) MgSO_4 has highest osmotic pressure
- (d) K_2SO_4 has highest osmotic pressure

Q22. In the extraction of metallic Zn from calamine ore (ZnCO_3) following steps are involved:

- (A) Distillation of impure Zn
- (B) Calcination of the ore
- (C) Hydraulic washing of ore
- (D) Reduction of ZnO using coke

Choose the **correct** order of steps from the options given below:

- (a) (C), (D), (B), (A)
- (b) (C), (B), (D), (A)
- (c) (A), (B), (C), (D)
- (d) (A), (C), (B), (D)

Q23. Match List - I with List - II.

	List - I		List - II
(A)	Glycosidic linkage	(I)	Monomeric units of nucleic acid
(B)	Peptide linkage	(II)	Sugar-heterocyclic base combination
(C)	Nucleotide	(III)	Linkage between two amino acid units

(D)	Nucleoside	(IV)	Linkage between two mono saccharide units
-----	------------	------	---

Choose the **correct** answer from the options given below:

- (a) (A)-(III), (B)-(IV), (C)-(I), (D)-(II)
 (b) (A)-(IV), (B)-(I), (C)-(II), (D)-(II)
 (c) (A)-(III), (B)-(IV), (C)-(II), (D)-(I)
 (d) (A)-(IV), (B)-(III), (C)-(I), (D)-(II)

Q24. Which property is observed in an emulsion?

- (a) Tyndall effect
 (b) Brownian motion
 (c) Tyndall effect and Brownian motion
 (d) Stable in nature

Q25. Which phenomenon occurs when an electric field is applied to a colloidal solution and electrophoresis is prevented?

- (A) Reverse osmosis takes place
 (B) Electroosmosis takes place
 (C) Dispersion medium begins to move in an electric field
 (D) Dispersion medium becomes stationary

Choose the **correct** answer from the options given below:

- (a) (A), (D) only
 (b) (B), (D) only
 (c) (A), (C) only
 (d) (B), (C) only

Q26. When electric current is passed through acidified water, 448 mL of hydrogen gas at STP is collected at the cathode in 965 seconds. The current passed in amperes is:

- (a) 1.0
 (b) 2.5
 (c) 3.0
 (d) 4.0

Q27. A square planar complex is formed by hybridization:

- (a) s, p_x, p_y, p_z
 (b) s, p_x, p_y, d_{z^2}
 (c) s, p_y, p_z, d_{xy}
 (d) $s, p_x, p_y, d_{x^2-y^2}$

Q28. The hormone responsible for preparing the uterus for implantation of fertilised egg is:

- (a) Progesterone
 (b) Testosterone
 (c) Epinephrine
 (d) Norepinephrine

Q29. Which of the following statement is **incorrect**?

- (a) Brownian movement stabilises **sol**.
 (b) Both emulsifier and peptising agents stabilises colloids but their actions are different.
 (c) Mixing two oppositely charged sols neutralizes their charges and stabilises the colloid.
 (d) The stability of lyophobic colloidal particles is due to preferential adsorption of ions on their surface from the solution.

Q30. Which of the following process is responsible for the formation of delta at a place where rivers meet the sea?

- (a) Emulsification
 (b) Colloid formation
 (c) Coagulation
 (d) Peptization

Q31. Match **List – I** with **List – II**.

	List – I		List – II
(A)	Ethanol and Diethyl ether	(I)	Lucas reagent
(B)	Methanol and Ethanol	(II)	Sodium metal
(C)	n-Butyl alcohol and tert. Butyl alcohol	(III)	Iodoform test
(D)	Phenol and Cyclohexanol	(IV)	Ferric Chloride

Choose the **correct** answer from the options given below:

- (a) (A)-(II), (B)-(IV), (C)-(III), (D)-(I)
 (b) (A)-(I), (B)-(II), (C)-(IV), (D)-(III)
 (c) (A)-(I), (B)-(IV), (C)-(III), (D)-(II)
 (d) (A)-(II), (B)-(III), (C)-(I), (D)-(IV)


Q32. o-Nitrophenol is less soluble in water than p-nitrophenol because:

- (a) o-Nitrophenol is more steam volatile than p-nitrophenol
 (b) o-Nitrophenol shows intramolecular H-bonding
 (c) o-Nitrophenol shows intermolecular H-bonding
 (d) Melting point of o-Nitrophenol is lower than p-nitrophenol

Q33. In the 3d series of transition elements, the element more likely to exhibit + 1 oxidation state is:

- (a) Fe
 (b) Mn
 (c) Cu
 (d) Ag

Q34. An aromatic compound 'A' of molecular formula C_7H_7ON undergoes a series of reaction as:



Identify compound D:

- (a) C_7H_5N
 (b) C_6H_6O
 (c) C_6H_7N
 (d) C_6H_6

Q35. The magnetic moment of Cu^{2+} ion ($Z = 29$) is:

- (a) $\sqrt{3}$
 (b) $\sqrt{8}$
 (c) $\sqrt{2}$
 (d) $\sqrt{4}$

Q36. The correct statements about Cr^{2+} and Mn^{3+} are: (atomic number of Cr = 24 and Mn = 25)

- (A) Cr^{2+} is a reducing agent
 (B) Mn^{3+} is an oxidising agent
 (C) Both Cr^{2+} and Mn^{3+} exhibit d^4 electronic configuration
 (D) When Cr^{2+} is used as reducing agent chromium ion attains d^5 electronic configuration

Choose the **correct** answer from the options given below:

- (a) (B), (D) and (A) only
- (b) (A), (B), (C) only
- (c) (C), (D), (A) only
- (d) (D), (A) only

Q37. The formation of micelle takes place only:

- (A) above a particular temperature called kraft temperature
- (B) above a particular concentration called critical micelle concentration
- (C) when catalyst is added

Choose the **correct** answer from the options given below:

- (a) (A), (B) and (C)
- (b) (B), (C) Only
- (c) (A), (C) Only
- (d) (A), (B) Only


Q38. Antiseptics and disinfectants either kill or prevent growth of micro-organism. Identify which of the following statements is **not** true?

- (A) Disinfectants harm the living tissues.
- (B) A 0.2% solution of phenol is an antiseptic while 1% solution acts as disinfectant.
- (C) Chlorine and Iodine are used as strong disinfectants.
- (D) Dilute solution of boric acid and hydrogen peroxide are antiseptics
- (E) Iodoform is also used as an antiseptic for wounds.

Choose the **correct** answer from the options given below:

- (a) (C) only
- (b) (D) only
- (c) (E) only
- (d) (A) and (B) only

Q39. In the following sequence of reactions:



The compound [C] is:

- (a) Butan-1-ol
- (b) Butan-2-ol
- (c) 1, 1-Dimethyl ethanol
- (d) 2-methyl propan-2-ol

Q40. An alkyl halide with molecular formula $\text{C}_6\text{H}_{13}\text{Br}$ on dehydrohalogenation gave two isomeric alkenes A and B with molecular formula C_6H_{12} . On reductive ozonolysis, A and B gave four compounds CH_3COCH_3 , CH_3CHO , $\text{CH}_3\text{CH}_2\text{CHO}$ and $(\text{CH}_3)_2\text{CHCHO}$. The alkyl halide is:

- (a) 2-Bromohexane
- (b) 2,2-Dimethyl-1-bromobutane
- (c) 2-Bromo-2,3-dimethylbutane
- (d) 3-Bromo-2-methylpentane

Direction (Q41-Q45) Answer the question on the basis of passage given below:

A ketone [A] which undergoes haloform reaction gives compound [B] on reduction. Compound [B] on heating with sulphuric acid gives compound [C] which forms monozonide [D]. The compound [D] on hydrolysis in the presence of zinc

dust gives compound [E]. Compound [E] on oxidation gives $\text{C}_2\text{H}_4\text{O}_2$ [F]

Q41. Identify Compound [A]

- (a) Butan-2-one
- (b) 3-methyl Pentan-2-one
- (c) Propanone
- (d) Acetophenone

Q42. The compound [B] formed can be prepared from reaction of aq.KOH. with:

- (a) $\text{CH}_3\text{CH}_2\text{CH}_2 - \text{Cl}$
- (b) $\text{CH}_3 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_3$
- (c) $\text{CH}_3 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_2 - \text{CH}_3$
- (d) $\text{CH}_3 - \underset{\text{Cl}}{\overset{\text{CH}_3}{\text{C}}} - \text{CH}_3$

Q43. Compound [E] in presence of dil NaOH gives:

- (a) But-2-enal
- (b) Methanol
- (c) 2-Methyl but-2-enal
- (d) Pent-2-enal

Q44. Compound [F] ($\text{C}_2\text{H}_4\text{O}_2$) on treatment with chlorine, in the presence of small amount of red phosphorus, gives:

- (a) $\text{CH}_3\text{CH}_2 - \text{Cl}$
- (b) CH_3COCl
- (c) ClCH_2COOH
- (d) $\text{ClCH}_2\text{CH}_2 - \text{Cl}$

Q45. The boiling point of [A] observed is exceptionally high because of:

- (a) Intermolecular-H-bond
- (b) Intramolecular-H-bond
- (c) Like interaction
- (d) Dipole-Dipole interaction

Direction (Q46-Q50) Answer the questions on basis of passage given below:

The noble gases He to Rn have close shell configuration and are monoatomic gases under normal conditions. Though mostly unreactive, but direct reaction of Xe with F_2 leads to a series of compounds with Xe oxidation numbers +2, +4 and +6. When certain compounds like phenol and quinol are crystallised in the presence of noble gas such as Ar, Kr and Xe a category of compound called clathrates are formed. In these the noble gas atoms are trapped in the cavities of the crystal lattices. The compounds which crystallise are known as hosts (H) while the noble gas atoms are called guests (G). The general formula of a clathrate compound is $n\text{HmG}$, where n is the number of host molecules and m is the number of atoms a molecule of guest.

Q46. In what molar ratio must Xe and F_2 be taken to give XeF_4 ?

- (a) 1 : 1
- (b) 1 : 2
- (c) 2 : 1
- (d) 1 : 5

Q47. XeF_4 and XeF_6 are:

- (a) reducing agent

- (b) oxidising agent
- (c) unreactive
- (d) strongly basic

Q48. Which force is responsible for holding Ar in its clathrate compound with quinol?

- (a) Van der Waal
- (b) Hydrogen bond
- (c) Ionic bond
- (d) Covalent bond

Q49. Match **List – I** with **List – II**.

	List – I		List - II
(A)	SF ₄	(I)	Bent 'T'
(B)	BrF ₃	(II)	Pyramidal
(C)	BrO ₃ ⁻	(III)	Seesaw
(D)	XeOF ₄	(IV)	Square pyramidal

Choose the **correct** answer from the options given below:

- (a) (A)-(III), (B)-(IV), (C)-(I), (D)-(II)
- (b) (A)-(IV), (B)-(II), (C)-(III), (D)-(I)
- (c) (A)-(III), (B)-(I), (C)-(II), (D)-(IV)
- (d) (A)-(I), (B)-(III), (C)-(II), (D)-(IV)

Q50. Which factors are responsible for making fluorine the strongest oxidising agent?

- (A) Highest electronegativity
- (B) High electron gain enthalphy
- (C) High bond dissociation enthalphy
- (D) Small size of fluorine

Choose the **correct** answer from the options given below:

- (a) (A), (B) and (D) only
- (b) (B), (C) and (D) only
- (c) (A), (C) and (D) only
- (d) (B) and (C) only